

Permanent Compared With Absorbable Suture in Apical Prolapse Surgery

A Systematic Review and Meta-analysis

Bracha L. Pollack, BA, Patrick Popiel, MD, Miriam C. Toaff, MD, Elizabeth Drugge, PhD, MPH, Adrienne Bielawski, BA, Ashley Sacks, BA, Moses Bibi, BS, Rebecca Friedman-Ciment, MS, Kira LeBron, MS, Leael Alishahian, BS, Dena Phillips, BA, Sara R. Rubino, BS, Sela Pollack, Rida S. Khan, BS, Eesha S. Khan, BS, Dominique Malacarne Pape, MD, and Cara L. Grimes, MD, MAS

OBJECTIVE: To explore how permanent compared with absorbable suture affects anatomic success in native tissue vaginal suspension (uterosacral ligament suspension and sacrospinous ligament suspension) and sacrocolpopexy with mesh.

DATA SOURCES: MEDLINE, EMBASE, and ClinicalTrials.gov were searched through March 29, 2022.

METHODS OF STUDY SELECTION: Our population included women undergoing apical prolapse surgery (uterosacral ligament suspension and sacrospinous ligament suspension and abdominal sacrocolpopexy). Our intervention was permanent suture for apical prolapse surgery, and our comparator was absorbable suture. We determined a single anatomic success proportion per study. Adverse events collected included suture and mesh exposure, surgery for suture and mesh complication, dyspareunia, and

granulation tissue. Abstracts were doubly screened, full-text articles were doubly screened, and accepted articles were doubly extracted. Quality of studies was assessed using GRADE (Grading of Recommendations Assessment, Development and Evaluation) criteria. In single-arm studies using either permanent or absorbable suture, random effects meta-analyses of pooled proportions were used to assess anatomic success. In comparative studies investigating both suture types, random effects meta-analyses of pooled risk ratios were used.

TABULATION, INTEGRATION, AND RESULTS: Of 4,658 abstracts screened, 398 full-text articles were assessed and 63 studies were included (24 vaginal suspension [13 uterosacral ligament suspension and 11 sacrospinous ligament suspension] and 39 sacrocolpopexy). At 2-year follow-up, there was no difference in permanent compared with absorbable suture in uterosacral ligament suspension and sacrospinous ligament suspension (proportional anatomic success rate 88% [95% CI 0.81–0.93] vs 88% [95% CI 0.82–0.92]). Similarly, at 18-month follow-up, there was no difference in permanent compared with absorbable suture in sacrocolpopexy (proportional anatomic success rate 92% [95% CI 0.88–0.95] vs 96% [95% CI 0.92–0.99]). On meta-analysis, there was no difference in relative risk (RR) of success for permanent compared with absorbable suture for uterosacral ligament suspension and sacrospinous ligament suspension (RR 1.11, 95% CI 0.93–1.33) or sacrocolpopexy (RR 1.00, 95% CI 0.98–1.03).

CONCLUSION: Success rates were similarly high for absorbable and permanent suture after uterosacral ligament suspension, sacrospinous ligament suspension, and sacrocolpopexy, with medium-term follow-up.

SYSTEMATIC REVIEW REGISTRATION: PROSPERO, CRD42021265848.

(Obstet Gynecol 2023;141:268–83)

DOI: 10.1097/AOG.0000000000005032

From the New York Medical College, and the Departments of Obstetrics and Gynecology, Urology, and Pharmacology, and the Department of Public Health, School of Health Sciences and Practice, New York Medical College, Hawthorne, New York.

Presented at the Society of Gynecologic Surgeons' 48th Annual Scientific Meeting, March 27–30, 2022, San Antonio, Texas; and at the AUGS/IUGA Scientific Meeting, June 14–18, 2022, Austin, Texas.

The authors thank Judy Moore for her help with administrative and research support.

Each author has confirmed compliance with the journal's requirements for authorship.

Corresponding author: Cara L. Grimes, MD, MAS, Departments of Obstetrics and Gynecology and Urology, New York Medical College, Hawthorne, NY; email: caragrimesmd@gmail.com.

Financial Disclosure:

Cara L. Grimes reports providing Expert testimony for Johnson and Johnson. The other authors did not report any potential conflicts of interest.

© 2023 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

ISSN: 0029-7844/23

Pelvic organ prolapse (POP) is common. One in eight women will undergo POP surgery in their lifetimes.¹ Abdominal sacrocolpopexy with mesh and native tissue vaginal suspensions, including uterosacral ligament suspension and sacrospinous ligament fixation, are commonly performed surgeries to treat apical POP.^{2–5}

There is conflicting evidence to support the choice of suture material during POP surgery and whether suture type affects anatomic or subjective outcomes after surgery. The choice of suture type often is determined by surgeon preference. Some studies have demonstrated that permanent suture has lower failure rates⁶ than absorbable suture, whereas others have shown similar anatomic outcomes.⁷ Permanent suture classically has been used at the vaginal apex and sacrum; however, use of permanent suture increases the risk of suture or mesh erosion into the vagina.^{5,8} Because of this, the use of delayed absorbable suture has gained popularity. Subsequently, a number of studies have shown similar rates of post-operative parameters such as POP recurrence and reoperation with the use of absorbable suture.^{8–10} A recent survey showed that surgeons preferred absorbable suture for uterosacral ligament suspension and sacrospinous ligament suspension¹¹ and that, although most surgeons used permanent sutures for the sacral attachment, most surgeons preferred absorbable suture for the vaginal attachment of sacrocolpopexy.^{8,11}

We aimed to systematically review the literature to explore the anatomic outcomes of using either permanent or absorbable suture for apical POP surgery. We hypothesized that permanent suture results in better anatomic outcomes but with more suture complications.

METHODS

No IRB approval was required for this work. PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines were followed. Registration with PROSPERO and full protocol can be found at: https://www.crd.york.ac.uk/PROSPERO/display_record.php?RecordID=265848.

SOURCES

We searched MEDLINE, EMBASE, and ClinicalTrials.gov from their inception through June 11, 2021, and then ran an updated search on March 29, 2022, and April 18, 2022 (for ClinicalTrials.gov). The search included numerous MeSH terms, such as “apical pelvic organ prolapse,” “sacrocolpopexy,” “utero-

sacral ligament suspension,” “sacrospinous ligament suspension,” “suture material,” and other associated text words (Appendix 1, available online at <http://links.lww.com/AOG/C974>). No filters or limits were used. Non-human studies and conference abstracts were excluded. Authors were not contacted for additional information.

STUDY SELECTION

Our population included studies of women who underwent apical prolapse surgery, including native tissue vaginal apical suspensions (uterosacral ligament suspension and sacrospinous ligament suspension) and sacrocolpopexy with type 1 polypropylene mesh. Studies of hysteropexy and McCall culdoplasty and those not including apical prolapse surgery were excluded. Additionally, we excluded studies that involved cadavers, tissue samples, and animal studies. Our intervention was defined as permanent suture (including polytetrafluoroethylene, polypropylene, poliglecaprone 25, silk, and polyamide). We chose to exclude braided polyester based on studies that have demonstrated high rates of adverse events and expert consensus that these sutures are not commonly used anymore.¹² Our comparator group was absorbable or delayed absorbable suture (polyglactin, polydioxanone, polyglyconate, glycolide, dioxanone, and trimethylene carbonate). No absorbable sutures were excluded from inclusion.

Our content experts (P.P., C.L.G., D.M.P.) collectively determined a single composite anatomic success proportion for each study. This determination took into account pelvic organ prolapse quantification (POP-Q) measurements, stage or grade, subjective symptoms of bulge, re-operations, and re-treatments. We were careful to make sure cases were not double counted and to arrive at a proportion that considered all of the above factors. Adverse events including suture exposure, surgery for suture or mesh complication, mesh erosion, dyspareunia, and granulation tissue were collected. Randomized controlled trials (RCTs), prospective or retrospective nonrandomized comparative and cohort studies, and single-arm case series were included. Video presentations, conference abstracts, and articles in languages other than English were excluded.

Seventeen reviewers independently screened abstracts and potentially relevant full-text articles in duplicate. Discrepancies were resolved by a third reviewer (P.P., C.L.G., D.M.P.). Abstracts were doubly screened using Abstrackr (<http://abstrackr.cebm.brown.edu>),¹³ then full-text articles were doubly screened. Data extraction was completed in duplicate

into customized forms by the same independent reviewers. Study and participant characteristics, intervention details, outcome definitions, results, and adverse events were extracted. The quality of studies was assessed using GRADE (Grading of Recommendations Assessment, Development and Evaluation) criteria.¹⁴ The risk of bias and methodologic quality of each study were assessed based on Cochrane risk of bias and other questions from the Newcastle-Ottawa Scale.^{15,16} Based on these potential biases, each study was graded by two to three reviewers as good (A), fair (B), or poor (C) quality.

In studies using either absorbable or permanent suture (single-arm studies), random effects meta-analyses of pooled proportions were used to assess anatomic success using Stata's Metaprop Statistical Program,¹⁷ which uses the Freeman-Tukey double arcsine transformation to stabilize the variances. In studies investigating both suture types (comparative studies), random effects model-restricted maximum likelihoods were used to meta-analyze risk ratios. The I^2 statistic was used to reflect heterogeneity; all subgroups were planned a priori, and studies reporting the same outcome were meta-analyzed regardless of the degree of heterogeneity. Adverse events were compared with the two-sample test of proportions. Analyses were performed using Stata 17, and $P < .05$ was considered statistically significant.

RESULTS

The literature search identified 4,658 abstracts; 398 full-text articles were retrieved and assessed in detail. In total, we included 63 studies that met eligibility criteria. We categorized the studies into 24 vaginal native tissue repair studies and 39 sacrocolpopexy studies (Fig. 1).

Twenty-four vaginal suspension studies with 2,309 patients met inclusion criteria. Two studies^{7,18} (both uterosacral ligament suspension) compared both suture types, and 22 used one suture type (12 permanent^{12,19–29} and 10 absorbable^{6,30–38}). Of these single-arm studies, 11 used uterosacral ligament suspension^{6,12,27–29,33–38} and 11 used sacrospinous ligament suspension.^{19–26,30–32} Of the patients undergoing uterosacral ligament suspension, 473 had absorbable suture and 1,044 had permanent suture. Of the patients undergoing sacrospinous ligament suspension, 609 had absorbable suture and 183 had permanent suture. The average age of the study populations ranged from 53 to 59 years, and average body mass index (BMI, calculated as weight in kilograms divided by height in meters squared) was between 22 and 30. Most studies reported POP-Q stage. When reported,

the majority of patients had at least stage 3 prolapse (Table 1).

We assigned two studies a quality grade of A, six were assigned a grade of B, and 16 were assigned a grade of C. Overall, four studies were RCTs, eight were comparative (one prospective, seven retrospective), and 12 were single-arm (six prospective, six retrospective). The mean follow-up time was 23 months (Table 1).

Of the 13 uterosacral ligament suspension studies, the majority placed sutures on both sides (11/13, 84.6%, two not reported) and the majority specifically referenced the Shull method or a modified Shull method (11/13, 84.6%). Most of the time, two sutures were placed in both uterosacral ligaments (7/13, 53.8%). Two studies (15%) did not report the number of sutures, and three studies (23.1%) placed three on each side. Suture caliber was reported most of the time (9/13, 69.2%), and most sutures were 2-0, 1-0, or 0 in caliber. Two studies (15%) of permanent suture actually included both absorbable and permanent sutures.

Of 11 sacrospinous ligament suspension studies, we found that sutures were placed solely at the right sacrospinous ligament 63.6% (7/11) of the time. One study placed sutures at the left ligament, one in both ligaments, and one at the right or in both ligaments; one study did not record suture attachment site(s). The majority of studies (8/11, 72.7%) described placing two sutures at each attachment site; three studies did not report suture number. Suture caliber was reported most of the time (8/11, 72.7%), with 0 or 1-0 used most often. Permanent suture (8/11, 72.7%) was used more often than absorbable suture (3/11, 27.3%) in the sacrospinous studies.

Overall, at 23-month follow-up, there was no difference in permanent compared with absorbable suture in uterosacral ligament suspension or sacrospinous ligament suspension (proportional anatomic success rate 88%, 95% CI 0.84–0.91). The proportional anatomic success rate of permanent suture in vaginal suspension ($n=14$) was 88% (95% CI 0.81–0.93); for absorbable suture in vaginal suspension ($n=12$), it was also 88% (95% CI 0.82–0.92) (Fig. 2, Appendix 2 [Appendix 2 is available online at <http://links.lww.com/AOG/C974>]). There was high heterogeneity between groups ($I^2=84.4\%$), with no difference in success rates across groups. Further subgroup analysis by type of vaginal suspension (uterosacral ligament suspension or sacrospinous ligament suspension) also showed no differences between suture types (Appendix 3, available online at <http://links.lww.com/AOG/C974>). The proportional anatomic success rate of permanent suture




Fig. 1. Selection process in accordance with PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-analyses) guidelines.

Pollack. Suture Type in Apical Prolapse Surgery. Obstet Gynecol 2023.

for uterosacral ligament suspension ($n=6$) was 83% (95% CI 0.68–0.94); for absorbable suture ($n=9$), it was 88% (95% CI 0.82–0.93). The proportional anatomic success rate of permanent suture for sacrospinous ligament suspension ($n=8$) was 91% (95% CI 0.85–0.95); for absorbable suture ($n=3$), it was 87% (95% CI 0.72–0.97).

Most studies (14/24, 58.3%) reported on adverse events, with eight reporting on presence or absence of suture exposure, six reporting on surgery for suture complications, seven reporting on dyspareunia, and

five reporting on granulation tissue. There was more suture exposure in the permanent suture group compared with the absorbable suture group, with a 5.6% difference (95% CI 0.03–0.09). There were more surgeries for suture and mesh in the permanent suture group compared with the absorbable suture group, with a 4.6% difference (95% CI 0.01–0.08). There was more granulation tissue in the permanent suture group compared with the absorbable suture group, with a 10.1% difference (95% CI 0.06–0.14). There

Table 1. Summary of Native Tissue Vaginal Suspension Studies

Study	Study Design (Quality)	No. of Participants	Age (y)	BMI (kg/m ²)	Preoperative Prolapse Severity	Mean Follow-up Time (mo)	Surgery Type (Suture Type)
Kowalski, 2021 ¹⁸	RCT (A)	44	P: 63.6±10.2 A: 62.1±13.7	P: 29.7±4.6 A: 28.4±6.2	POP-Q stage 3–4: P: 68% A: 64%	12	Uterosacral, Shull method (P: polytetrafluoroethylene; A: polydioxanone)
Bradley, 2018 ⁷	Comparative, retrospective (B)	242	P: 64.8±8.1 A: 62.2±10.5	P: 28.5±5.4 A: 27.1±4.9	POP-Q stage 3–4: P: 79% A: 59%	12	Uterosacral, Shull method (P: surgeon preference; A: polyglyconate)
Da Silva, 2020 ¹⁹	RCT (B)	59	65.4±8	27.9±3.8	POP-Q mean: Ba: +3.76 C: +1.46 Bp: +1.81	60	Sacrospinous (polypropylene)
Goldberg, 2001 ²⁰	Comparative, retrospective (C)	168	67 (30–89) [†]	NR	POP-Q stage 3–4: greater than 82%	39	Sacrospinous (polytetrafluoroethylene)
Mothes, 2015 ²¹	Single group, prospective (C)	110	63 (39–89) [†]	NR	POP-Q stage 3–4: 100%	14	Sacrospinous (polyamide)
Maggiore, 2013 ²²	Comparative, prospective (C)	86	61.1	24.6	POP-Q stage 3–4: 79%	36	Sacrospinous (polypropylene)
Peng, 2010 ²³	Single group, retrospective (C)	35	60.2	NR	POP-Q mean: Ba: +0.82 Bp: –1.97	13	Sacrospinous (silk)
Astepe, 2019 ²⁴	Comparative, retrospective (C)	43	59±8.3	30±3.5	POP-Q stage 2–4: 100%	17	Sacrospinous (permanent, not specified)
Salvat, 1996 ²⁵	Single group, retrospective (C)	20	64.7 (46–86) [†]	NR	NR	72	Sacrospinous (polyamide)
Shastri, 2020 ²⁶	Single group, retrospective (C)	106	53 (40–70) [†]	NR	POP-Q stage 3–4: 100%	6	Sacrospinous (polypropylene)
Wheeler, 2007 ²⁷	Single group, retrospective (C)	35	64.2±10.2	27±4.5	POP-Q mean: C: –1.5 Ba: +2.2 Bp: +0.21	24	Uterosacral, not Shull (polypropylene)
Lee, 2021 ¹²	Comparative, retrospective (C)	149	67 (56–78) [†]	24.8±2.6	POP-Q stage 3–4: 76%	12	Uterosacral, Shull method (polypropylene and polydioxanone)
Nager, 2021 ²⁸	RCT (A)	87	66.2±7.4	28.1±4.4	POP-Q stage 3–4: 78%	60	Uterosacral, Shull method (1 polypropylene and 1 absorbable on each side)
Barber, 2000 ²⁹	Single group, retrospective (C)	41	65±11	NR	POP-Q stage 3–4: 87%	15.5	Uterosacral, Shull method (1 permanent and 1 delayed absorbable on each side)
Mowat, 2018 ³⁰	Single group, prospective (C)	51	66.1±8.9	27.6±3.9	POP-Q median: C: –2 Ba: +1 Bp: 0	17	Sacrospinous (polydioxanone)
Dangal, 2018 ³¹	Single group, prospective (C)	95	NR	NR	POP-Q stage 3–4: 87%	6	Sacrospinous (polydioxanone)
Greisen, 2021 ³²	Single group, prospective (C)	103	65 [‡]	26 [‡]	POP-Q stage 3–4: 16.5%	6	Sacrospinous (polydioxanone)
Chung, 2012 ⁶	Comparative, retrospective (B)	141	60.1 (1)	28±0.5	POP-Q stage 3–4: 63%	5	Uterosacral, Shull method (polydioxanone)
Jeffery, 2009 ³³	Single group, retrospective (C)	53	63±11.6	28±6.3	BW grade 2–4: 66%	15	Uterosacral, modified high uterosacral ligament suspension (polydioxanone)
Wong, 2011 ³⁴	Single group, retrospective (C)	57	56±6.9	30±5.4	POP-Q median: C: –2 Ba: +1 Bp: –1	12	Uterosacral, Shull method (polyglyconate)
Spelzini, 2017 ³⁵	RCT (B)	124	56.7±9	24.5±3.5	POP-Q median: C: 0 Ba: +1 Bp: –1	12	Uterosacral, Shull method (polydioxanone)
Schiavi, 2017 ³⁶	Single group, retrospective (C)	146	61.6±8.3	27.3±3.8	POP-Q mean: C: +2.3 Ba: +3.2 Bp: +1.5	48	Uterosacral, Shull method (polydioxanone)

(continued)

Table 1. Summary of Native Tissue Vaginal Suspension Studies (continued)

Study	Study Design (Quality)	No. of Participants	Age (y)	BMI (kg/m ²)	Preoperative Prolapse Severity	Mean Follow-up Time (mo)	Surgery Type (Suture Type)	
Rappa, 2016 ³⁷	Comparative, retrospective (B)	360	Normal weight: 69.1 (5.7) Overweight or obesity: 69.5 (11.4)	Normal weight: 22.1±1.2 Overweight or obesity: 30.7 mean (4.4)	POP-Q stage 3–4: Normal weight: 65% Overweight or obesity: 65.9%	12.5	Uterosacral, Shull method (polydioxanone)	
Chill, 2021 ³⁸	Comparative, retrospective (B)	112	63.2±8.5	27.2±4.1	POP-Q stage 3–4: 95.5%	19.4	Uterosacral (polydioxanone)	
Study	No. of Sutures	Side of Attachment	Suture Caliber	Definition of Treatment Success*	Anatomic Success [n/N (%)]		Adverse Events	
					Permanent Suture	Absorbable Suture	Permanent Suture	Absorbable Suture
Kowalski, 2021 ¹⁸	4 (2/side)	Both	CV-2, 0	1–4	15/20 (75)	19/20 (90)	2/20 suture exposure 2/20 dyspareunia 0/22 granulation tissue	1/20 suture exposure 1/20 dyspareunia 0/22 granulation tissue
Bradley, 2018 ⁷	NR	Both	NR	1, 3, 4	43/54 (80)	156/188 (83)	1/53 suture exposure 5/53 surgery for suture complication (pain) 3/53 granulation tissue	0/184 suture exposure 6/184 surgery for suture complication (pain) 5/184 granulation tissue
Da Silveira, 2020 ¹⁹	NR	Right	0	1, 3, 4	45/59 (76)	NA	1/59 suture exposure	NA
Goldberg, 2001 ²⁰	2	Right	0	1, 4	151/168 (90)	NA	4/168 dyspareunia	NA
Mothes, 2015 ²¹	2	Both	0	1–4	104/110 (95)	NA	NR	NA
Maggiore, 2013 ²²	2	Right	NR	1, 2	75/86 (87)	NA	10/86 dyspareunia	NA
Peng, 2010 ²³	2	Right	No. 7	1, 2	33/35 (94)	NA	2/35 suture exposure 2/35 surgery for suture complication	NA
Astepe, 2019 ²⁴	2	Right	NR	1–3	37/43 (86)	NA	NR	NA
Salvat, 1996 ²⁵	NR	Right or both	“Decimal 4”	1	18/20 (90)	NA	NR	NA
Shastri, 2020 ²⁶	2	Left	1	1	87/88 (99)	NA	NR	NA
Wheeler, 2007 ²⁷	4 (2/side)	Both	0 or 1	1–4	26/26 (100)	NA	NR	NA
Lee, 2021 ¹²	6 (3/side) 1 absorbable and 2 permanent on each side	Both	0 polypropylene and 1-0 polydioxanone	1–4	127/149 (85)	NA	7/149 suture exposure 9/149 surgery for suture complication 25/149 granulation tissue (surgery)	NA
Nager, 2021 ²⁸	4 (2/side)	Both	0 or 2-0	1, 2, 4	42/78 (54)	NA	19/91 suture exposure 2/86 dyspareunia 11/91 granulation tissue	NA
Barber, 2000 ²⁹	4 (2/side)	Both	1-0 for both	1, 2	37/41 (90)	NA	NR	NA
Mowat, 2018 ³⁰	2	Right	NR	1, 2	NA	41/43 (96)	NA	1/51 surgery for suture complication 12/48 dyspareunia

(continued)

Table 1. Summary of Native Tissue Vaginal Suspension Studies (continued)

Study	No. of Sutures	Side of Attachment	Suture Caliber	Definition of Treatment Success*	Anatomic Success [n/N (%)]		Adverse Events	
					Permanent Suture	Absorbable Suture	Permanent Suture	Absorbable Suture
Dangal, 2018 ³¹	NR	Right	1	1, 2	NA	32/38 (84)	NA	3/95 surgery for suture complication
Greisen, 2021 ³²	2	NR	0	1, 2	NA	77/102 (75)	NA	NR
Chung, 2012 ⁶	2–4/side	1 or both sides	NR	1	NA	133/141 (94)	NA	7/141 suture exposure 0/141 surgery for suture complication
Jeffery, 2009 ³³	3 on each side	Both	0	1, 2	NA	47/53 (89)	NA	NR
Wong, 2011 ³⁴	2 on each side	Both	0	1, 2	NA	56/57 (98)	NA	1/57 suture exposure
Spelzini, 2017 ³⁵	3	Both	0	1–3	NA	63/73 (86)	NA	5/73 dyspareunia
Schiavi, 2017 ³⁶	NR	Both	0	1, 2	NA	133/146 (92)	NA	2/146 dyspareunia 3/146 granulation tissue
Rappa, 2016 ³⁷	4 (2/side)	Both	NR	1, 3, 4	NA	289/360 (80)	NA	NR
Chill, 2021 ³⁸	4 (2/side)	Both	2-0	1–3	NA	78/111 (70)	NA	NR

BMI, body mass index; RCT, randomized controlled trial; P: permanent; A: absorbable; POP-Q, pelvic organ prolapse quantification; NR: not reported; NA: not applicable.

Data are mean±SD unless otherwise specified.

* A single composite anatomic success proportion was determined by our content experts and was based on: 1) anatomic success (POP-Q stage, or grade), 2) subjective symptoms of bulge, 3) re-operations, and 4) re-treatments.

† Mean (range)

* Median (range).

was no difference in dyspareunia (Appendix 4, available online at <http://links.lww.com/AOG/C974>).

Thirty-nine studies with 3,349 patients met inclusion criteria for the sacrocolpopexy category. Three compared both suture types,^{9,39,40} and 36 used one suture type (20 permanent^{41–60} and 16 absorbable^{8,61–75}); 2,045 patients underwent sacrocolpopexy with permanent suture, and 1,304 with absorbable suture. Overall, nine studies were RCTs, 12 were comparative (two prospective, nine retrospective, one unclear direction), and 18 were single-arm (six prospective, 12 retrospective). The average age of the study population ranged from 43 to 70 years, and average BMI was between 20 and 29. Most studies reported POP-Q stage. When reported, the majority of patients had at least stage 3 prolapse. Several studies reported mean POP-Q measurements. We assigned four studies a quality grade of A, 15 were assigned a grade of B, and 20 were assigned a grade of C. The mean follow-up was 18.2 months (Table 2).

Suture caliber was reported in 21 of 39 (53.8%) studies; there was diversity in suture choice, including 1-0, 2-0, 3-0, CV-2, and CV4. Nineteen studies reported suture number; this varied widely, with a range of 3–12 interrupted sutures placed on each side of the vaginal mesh attachment. Two studies also re-

ported using one running suture. When placing sacral attachment sutures, most studies described using permanent suture (32/39, 82.0%).

Overall, at 18 months follow-up, there was no difference in permanent suture compared with absorbable suture in sacrocolpopexy (proportional anatomic success rate 94%, 95% CI 0.91–0.96). The proportional anatomic success rate of permanent suture for sacrocolpopexy (n=23) was 92% (95% CI 0.88–0.95); for absorbable suture for sacrocolpopexy (n=19), it was 96% (95% CI 0.92–0.99). There was considerable heterogeneity between groups ($I^2=87.2\%$), with no difference in success rates across groups (Fig. 3) (Appendix 5, available online at <http://links.lww.com/AOG/C974>). On meta-analysis, there was no difference in relative risk (RR) of success compared with failure for permanent compared with absorbable suture for sacrocolpopexy (n=3) (RR 1.00, 95% CI 0.98–1.03), with low heterogeneity ($I^2=0.03\%$). (Fig. 4)

Adverse events were reported in 33 of the 39 sacrocolpopexy studies (84.6%), with 12 studies reporting on the presence or absence of suture exposure, six studies reporting on surgery for suture and mesh complications, 28 studies reporting on mesh erosion, six studies reporting on dyspareunia, and no




Fig. 2. Proportional outcome analysis of vaginal suspension surgeries (including uterosacral and sacrospinous) by suture type (permanent vs absorbable). ES, effect size.

Pollack. Suture Type in Apical Prolapse Surgery. *Obstet Gynecol* 2023.

studies reporting on granulation tissue. There were more suture exposures in the permanent suture group compared with the absorbable suture group, with a 2.7% difference (95% CI 0.01–0.04). Permanent suture was 5.26 times more likely to become exposed than absorbable suture (RR 5.26, 95% CI 1.56–17.69) (Appendices 6 and 7, available online at <http://links.lww.com/AOG/C974>). There were more surgeries for suture and mesh complications in the permanent suture group compared with the absorbable suture group, with a 1.7% difference (95% CI 0.01–0.03). There was no difference in mesh erosion or dyspareunia between permanent and absorbable suture.

DISCUSSION

The choice of suture used in the surgical repair of apical prolapse traditionally has been rooted in

institutional and personal preferences, not driven by data. A growing body of research has been developing to more appropriately guide these clinical decisions. Although intuitively many surgeons thought permanent suture would offer a more durable and long-lasting repair, and thus higher success rates, there is concern that there may be more adverse events associated with permanent suture. Interestingly, our data found no difference in anatomic outcomes between permanent and absorbable suture at moderate-length follow-up. We find this very reassuring because it supports the use of absorbable suture to minimize adverse events such as suture exposure, granulation tissue, and repeat surgery for suture or mesh complications, without sacrificing success.

Our data support the growing body of literature that absorbable suture is a safe and effective option for

Table 2. Summary of Sacrocolpopexy Studies

Study	Study Design (Quality)	Suture Type	No. of Participants	Age (y)	BMI (kg/m ²)	Preoperative Prolapse Severity	Follow-up Time (mo)
Powell, 2021 ⁹	Comparative, retrospective (C)	Polypropylene and polydioxanone	119	P: 62.1±13.4 A: 63.9±10	P: 28.8±6.8 A: 27.7±5.2	POP-Q mean P: C: -3.1 Ba: +2.1 Bp: +0.8 A: C: -3.7 Ba: +2.4 Bp: -0.2	P: 23 A: 10.4
Matthews, 2020 ³⁹	RCT (A)	Polydioxanone and polytetrafluoroethylene	182	59±10	27.5±5	POP-Q stage 3–4: 72%	12
Tan-Kim, 2014 ⁴⁰	Comparative, retrospective (B)	Polydioxanone and polypropylene	193	P: 61±9 A: 60±9	NR	POP-Q median stage: 3	18
Culligan, 2013 ⁴¹	RCT (B)	Polytetrafluoroethylene	58	56.2±8.5	25.6±3.6	POP-Q mean: C: -0.8 Ba: +1.8 Bp: -0.3	12
Elliot, 2006 ⁴²	Single group, prospective (B)	Polytetrafluoroethylene	21	67 (47–83) [†]	NR	POP-Q stage 3–4: 100%	24
Ross, 2005 ⁴³	Single group, prospective (B)	Polypropylene	51	67 (41–83) [†]	NR	POP-Q stage 3–4: 85%	12
Davila, 2020 ⁴⁴	Comparative, retrospective (C)	Permanent (type not specified)	30	70	29	POP-Q stage 3–4: Ultrasound-assisted: 73% Standard: 80%	6
Culligan, 2005 ⁴⁵	RCT (A)	Polytetrafluoroethylene	45	60.4±10.1	28.4±4.7	POP-Q mean: C: -2.3 Ba: +0.9 Bp: +0.5	12
Gracia, 2015 ⁴⁶	Comparative, prospective (B)	Polypropylene	30	43.5±7.6	24.3±3.6	POP-Q stage 3–4: 62.6%	6
Nosti, 2016 ⁴⁷	Comparative, retrospective (B)	Polytetrafluoroethylene	104	54.8 [†]	26.2 [†]	POP-Q stage 3–4: 65.9%	12
Benson, 2010 ⁴⁸	Comparative, retrospective (C)	Polyglactin	33	62.8 [†]	NR	POP-Q stage 3–4: 100%	29 1,365
Culligan, 2020 ⁴⁹	Single group, prospective (B)	Polytetrafluoroethylene	253	57.7±9.4	26±4.1	POP-Q stage 3–4: 59.5%	66
Salamon, 2013 ⁵⁰	Single group, retrospective (C)	Polytetrafluoroethylene	64	54.4 (35–76) [†]	25.6 (19–35.8) [†]	POP-Q stage 3–4: 62.5%	12
Salamon, 2013 ⁵¹	Single group, prospective (B)	Polytetrafluoroethylene	118	56.63±7.85	26.02±3.98	POP-Q stage 3–4: 56%	12
Tate, 2011 ⁵²	RCT (A)	Polytetrafluoroethylene	29	58±9	NR	NR	60
Martin, 2015 ⁵³	Comparative, retrospective (C)	Polytetrafluoroethylene	181	55.8±9.7	27.4±4.8	POP-Q stage median: 3 (1–4)	3
Sato, 2021 ⁵⁴	Single group, retrospective (C)	P (type not specified)	46	70.0±7.4	23.6±3	POP-Q stage 3–4: 32.6%	12
Kenton, 2016 ⁵⁵	Comparative, prospective (B)	Polytetrafluoroethylene	66	59.5±9.9	27.7±5.7	POP-Q mean: C: +0.5 Ba: +2.5 Bp: -0.2	12
Belsante, 2013 ⁵⁶	Single group, retrospective (C)	Polyglactin	35	65 (37–79) [†]	24.6 (18–30) [†]	POP-Q mean: C: -1.1	28
Elliot, 2007 ⁵⁷	Single group, prospective (C)	Polytetrafluoroethylene	35	67 (47–83) [†]	NR	POP-Q stage 3–4: 100%	36

(continued)

Table 2. Summary of Sacrocolpopexy Studies (continued)

Study	Study Design (Quality)	Suture Type	No. of Participants	Age (y)	BMI (kg/m ²)	Preoperative Prolapse Severity	Follow-up Time (mo)
Mueller, 2016 ⁵⁸	Single group, retrospective (C)	Polytetrafluoroethylene	352	57.2±8.5	26.2±7.8	POP-Q stage 3–4: 80%	3
Van den Akker, 2019 ⁵⁹	Single group, retrospective (C)	P (type not specified) plus staples	178	Median: 66 (59–71) [§]	Median: 26.4 (23.4–28.7) [§]	POP-Q stage 3–4: 31%	35
Giannini, 2022 ⁶⁰	Single group, retrospective (C)	Polypropylene	60	62.2±7	24.8±2	POP-Q stage 2–4: 69%	24
Bazzi, 2019 ⁶¹	Comparative, retrospective (C)	A (type not specified)	131	64.5±10.1	28.5±5.2	POP-Q stage 3–4: 82%	12
Morciano, 2018 ⁶²	RCT (B)	Polydioxanone	84	65 (58–72) [§]	25 (22–27) [§]	POP-Q median stage: 3 (2–4)	12
Tagliaferri, 2021 ⁸	RCT (A)	Polyglactin	150	59±8	26.4±4.1	POP-Q median: C: +3 Ba: +3 Bp: +1	12
Berger, 2020 ⁶³	RCT (B)	A (type not specified)	46	60.7±8.8	27.5±4.6	POP-Q stage 3–4: 84.9%	12
Cvach, 2012 ⁶⁴	Comparative, unclear direction (C)	Polydioxanone	9	50 (37–66) [§]	25.5 (21–29) [†]	POP-Q stage 3–4: 100%	19
Tan-Kim, 2015 ⁶⁵	RCT (B)	Polydioxanone	64	58.7±9.4	25.6±3.9	POP-Q stage median: 3 (2–4)	12
Gillera, 2008 ⁶⁶	Single group, retrospective (C)	A (type not specified)	29	64±11	26±3.4	NR	23
Liu, 2020 ⁶⁷	Single group, retrospective (C)	Glycolide, dioxanone, and trimethylene carbonate (V-Loc 90)	49	60.8±9.3	23.7±3.2	POP-Q stage 3–4: 94.8%	3
Borohay, 2014 ⁶⁸	Single group, retrospective (C)	Barbed delayed absorbable (V-Loc 180)	20	54.3±11.4	29.1±4.7	POP-Q stage 3–4: 75%	17.3
Shepherd, 2010 ⁶⁹	Comparative, retrospective (B)	Polydioxanone	415	60.7	NR	POP-Q stage 3–4: 69.2%	1.5
Stubbs, 2011 ⁷⁰	Single group, retrospective (C)	Polydioxanone	36	70	27	POP-Q stage 3–4: 70%	3.8
Shekhar, 2020 ⁷¹	Single group, prospective (C)	Delayed absorbable	20	42.3±15.9	24.7±3.3	POP-Q mean: C: +2 Ba: 0 Bp: –3	16
Balsamo, 2018 ⁷²	Comparative, retrospective (B)	Polyglycolic	73	69±10	20.9 (15.6–32.7) [§]	POP-Q stage 3–4: 86.3%	94
Kallidonis, 2017 ⁷³	Single group, retrospective (C)	Trimethylene carbonate (V loc)	20	63 (50–79) [†]	26.7 (22.1–31.3) [†]	POP-Q stage 3–4: 25%	13.6
Liu, 2018 ⁷⁴	Single group, retrospective (C)	Trimethylene carbonate (V loc)	15	63.9 (39–80) [†]	27 (18.4–32.4) [†]	POP-Q stage 3–4: 60%	3
Reisenauer, 2022 ⁷⁵	RCT (B)	Polyglycolic acid	195	65.1±9.2	25.5±3.3	POP-Q stage 3–4: 30%	6

Study	Sacral Suture Type	No. of Sutures on Vaginal Mesh Attachment	Suture Caliber	Definition of Treatment Success*	Anatomic Success [n/N (%)]		Adverse Events	
					Permanent Suture	Absorbable Suture	Permanent Suture	Absorbable Suture
Powell, 2021 ⁹	NR	NR	NR	1	49/49 (100)	70/70 (100)	9/49 suture exposure 2/49 surgery for suture complication	2/70 suture exposure 0/70 surgery for suture complication

(continued)

Table 2. Summary of Sacrocolpopexy Studies (continued)

Study	Sacral Suture Type	No. of Sutures on Vaginal Mesh Attachment	Suture Caliber	Definition of Treatment Success*	Anatomic Success [n/N (%)]		Adverse Events	
					Permanent Suture	Absorbable Suture	Permanent Suture	Absorbable Suture
Matthews, 2020 ³⁹	Permanent	4 or more on each side	2-0 (both)	1,2,4	88/95 (93)	83/87 (95)	2/95 suture exposure 1/95 surgery for suture complication 5/95 mesh erosion	0/87 suture exposure 0/87 surgery for suture complication 7/87 mesh erosion
Tan-Kim, 2014 ⁴⁰	NR	Ant: 6 or more Post: 6 or more	2-0	1	143/148 (97)	43/45 (96)	4/148 suture exposure 25/148 mesh erosion	0/45 suture exposure 6/45 mesh erosion
Culligan, 2013 ⁴¹	Permanent	Ant: 6-10/side	NR	1	50/58 (86)	NA	0/58 suture exposure 3/58 dyspareunia	NA
Elliot, 2006 ⁴²	Permanent	NR	1-0	1	18/20 (90)	NA	2/20 mesh erosion	NA
Ross, 2005 ⁴³	Permanent	Ant: 5-6 Post: 3-5	NR	1	43/51 (84)	NA	4/51 mesh erosion 4/51 dyspareunia	NA
Davila, 2020 ⁴⁴	Permanent	NR	NR	1 and 4	29/30 (97)	NA	NR	NA
Culligan, 2005 ⁴⁵	Permanent	Ant: 9-5 Post: 6-12	NR	1	41/45 (91)	NA	2/45 mesh erosion	NA
Gracia, 2015 ⁴⁶	Permanent	NR	NR	1	21/30 (70)	NA	0/30 mesh erosion 0/30 dyspareunia	NA
Nosti, 2016 ⁴⁷	Permanent	Ant: 4-6 Post: 4-6	CV-2	1 and 2	93/104 (89)	NA	2/104 suture exposure 4/104 mesh erosion	NA
Benson, 2010 ⁴⁸	Permanent	Ant: 6-8 Post: 6-8	2-0	3	31/33 (94)	NA	0/33 mesh erosion 0/33 dyspareunia	NA
Culligan, 2020 ⁴⁹	Permanent	NR	CV-4	1 and 2	226/253 (89)	NA	0/253 mesh erosion	NA
Salamon, 2013 ⁵⁰	Permanent	NR	CV-4	1	57/64 (89)	NA	1/64 mesh erosion	NA
Salamon, 2013 ⁵¹	Permanent	NR	CV-4	1 and 2	105/118 (89)	NA	0/118 mesh erosion 2/118 dyspareunia	NA
Tate, 2011 ⁵²	Permanent	NR	CV-4	1-3	26/29 (90)	NA	3/29 mesh erosion	NA
Martin, 2015 ⁵³	Permanent	NR	NR	3,4	160/165 (97)	NA	5/165 suture exposure 2/165 surgery for suture complication	NA
Sato, 2021 ⁵⁴	Permanent	5 on each side	Ant: 3-0 Post: 2-0	1	43/46 (93)	NA	NR	NA
Kenton, 2016 ⁵⁵	Permanent	NR	NR	1 and 2	65/66 (98)	NA	1/66 suture exposure 0/66 mesh erosion	NA
Belsante, 2013 ⁵⁶	Permanent	NR	NR	1	35/35 (100)	NA	1/35 mesh erosion	NA
Elliot, 2007 ⁵⁷	Permanent	NR	NR	1	34/35 (97)	NA	0/35 suture exposure 0/35 surgery for suture complication 2/35 mesh erosion	NA
Mueller, 2016 ⁵⁸	Permanent	6-10 on both sides	CV-2	1	312/352 (89)	NA	5/458 mesh erosion	NA
Van den Akker, 2019 ⁵⁹	Tacks or sutures (type NR)	NR	NR	1	62/118 (59)	NA	4/178 surgery for mesh complication	NA
Giannini, 2022 ⁶⁰	Permanent	Ant: 6 Post: 6	NR	1	57/60 (95)	NA	0/60 mesh erosion	NA
Bazzi, 2019 ⁶¹	NR	NR	NR	1	NA	99/125 (79)	NA	6/125 mesh erosion

(continued)

Table 2. Summary of Sacrocolpopexy Studies (continued)

Study	Sacral Suture Type	No. of Sutures on Vaginal Mesh Attachment	Suture Caliber	Definition of Treatment Success*	Anatomic Success [n/N (%)]		Adverse Events	
					Permanent Suture	Absorbable Suture	Permanent Suture	Absorbable Suture
Morciano, 2018 ⁶²	A	Ant: 6 Post: 5 or running locking	3–0	1	NA	82/84 (98)	NA	NR
Tagliaferri, 2021 ⁸	Permanent	Ant: 5–10 Post: 4–6	2–0	1	NA	75/75 (100)	NA	0/75 mesh erosion
Berger, 2020 ⁶³	Permanent	Minimum 4/side	NR	1, 2, 3	NA	41/46 (89)	NA	NR
Cvach, 2012 ⁶⁴	Permanent	NR	2–0	1	NA	5/8 (63)	NA	2/8 mesh erosion
Tan-Kim, 2015 ⁶⁵	Permanent	Ant: 6–8 Post: 6–8	NR	1	NA	50/55 (91)	NA	2/55 mesh erosion
Gillera, 2008 ⁶⁶	Permanent	NR	NR	1,3	NA	29/29 (100)	NA	1/29 suture exposure 1/29 dyspareunia 3/49 mesh erosion
Liu, 2020 ⁶⁷	Permanent	“2–3 rows horizontally”	2–0	1,2	NA	49/49 (100)	NA	3/49 mesh erosion
Borohay, 2014 ⁶⁸	Permanent	NA	3–0	1	NA	20/20 (100)	NA	0/20 suture exposure 0/20 mesh erosion
Shepherd, 2010 ⁶⁹	1 absorbable	Ant: 3 Post: 6	2–0	1	NA	254/254 (100)	NA	0/254 suture exposure 2/254 mesh erosion
Stubbs, 2011 ⁷⁰	Permanent	1 running suture on each side	1–0	1	NA	33/36 (92)	NA	NR
Shekhar, 2020 ⁷¹	Permanent	4–6 sutures on each side	NR	1	NA	19/20 (95)	NA	0/20 mesh erosion
Balsamo, 2018 ⁷²	A	Ant: 4 Post: 4	1–0	1	NA	67/73 (92)	NA	1/73 mesh erosion
Kallidonis, 2017 ⁷³	A or permanent (titanium tack)	1 running suture on each side	3–0	1	NA	16/20 (80)	NA	0/20 mesh erosion
Liu, 2018 ⁷⁴	Permanent	NR	2–0	1	NA	15/15 (100)	NA	NR
Reisenauer, 2022 ⁷⁵	Permanent	NR	NR	1	NA	94/96 (98)	NA	0/96 suture exposure 0/96 surgery for suture complication 0/96 mesh erosion

BMI, body mass index; P: permanent; A: absorbable; POP-Q, pelvic organ prolapse quantification; RCT, randomized controlled trial; NR: not reported; Ant, anterior; Post, posterior; NA: not applicable.

Data are mean±SD unless otherwise specified.

* A single composite anatomic success proportion was determined by our content experts and was based on: 1) anatomic success (POP-Q stage, or grade), 2) subjective symptoms of bulge, 3) re-operations, and 4) re-treatments.

† Mean (range)

‡ No SD because two groups combined.

§ Median (range).

POP repair. For native tissue repair, a recent RCT found that absorbable suture was not inferior to permanent suture for uterosacral ligament suspension when comparing point C on the POP-Q 12 months postoperatively.¹⁸ In a 2020 systematic review looking at surgical success rates and mesh-related complications for absorbable suture compared with permanent suture in uterosacral ligament suspension, Peng et al found similar surgical success and failure rates. Furthermore, they found lower rates of suture exposure and erosion and need for suture removal with absorbable suture.⁷⁶ The existing data regarding

suture choice for sacrocolpopexy have been similar. In a recent randomized noninferiority trial, Tagliaferri et al found that, at 12 months postoperatively, delayed absorbable suture was noninferior to permanent suture when used for vaginal mesh attachment.⁸ Matthews et al³⁹ found no difference in suture and mesh erosion when comparing delayed absorbable suture and permanent suture in a large, multicenter RCT.

Our systematic review was performed using rigorous extraction methodology, and a composite definition of anatomic success verified by three experts was used when evaluating studies, despite significant

Downloaded from http://journals.lww.com/igreenjournal by 13Mps9n+0V11Mbe5QsRLc6ZUJOMK/rvXh4Th8pGCM
+UW3XJtIjboV01daEjBomFzeZv3MteioX+D+r1qT9g4Uy2+3N1YINNNXHHVZ2YwXKOYEardobd0i4DQouUH2U3SR0bFRZVW
96C6SizwFzyHXOF on 05/01/2023




Fig. 3. Proportional outcome analysis of sacrocolpopexy vaginal mesh attachment by suture type (permanent vs absorbable). ES, effect size.

Pollack. Suture Type in Apical Prolapse Surgery. *Obstet Gynecol* 2023.




Fig. 4. Forest plot of sacrocolpopexy vaginal mesh attachment by suture type (permanent vs absorbable). REML, restricted maximum likelihood.

Pollack. Suture Type in Apical Prolapse Surgery. *Obstet Gynecol* 2023.

heterogeneity in the study populations. Further, we had a mean follow-up of 1–2 years. It is important to note a number of limitations. The majority of the studies included were level B and C evidence. Additionally, considering the limited data that exist on this topic, we used studies with significant heterogeneity. The studies included represent a wide age range and diverse populations. Although this heterogeneity may actually make these data more applicable to a wider scope of patients that we encounter, this may also account for differences in the outcomes. Finally, a benefit of meta-analysis is that more patients are factored into the analysis, but our findings for sacrocolpopexy (Fig. 4, Appendix 7 [Appendix 7 is available online at <http://links.lww.com/AOG/C974>]) are primarily driven by the Powell study (weight 68.73).⁹

Our data should reassure surgeons using absorbable suture for apical prolapse surgery, specifically vaginal suspensions, including uterosacral ligament suspension and sacrospinous ligament suspension and sacrocolpopexy. Absorbable suture and permanent suture (excluding braided polyester) have similar success rates when used for apical prolapse repair. Further, it is likely that absorbable suture will lead to fewer suture exposures and complications and fewer repeat surgeries for suture complications.

REFERENCES

- Wu JM, Matthews CA, Conover MM, Pate V, Jonsson Funk M. Lifetime risk of stress urinary incontinence or pelvic organ prolapse surgery. *Obstet Gynecol* 2014;123:1201–6. doi: 10.1097/AOG.0000000000000286
- Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, Ulmsten U, et al. The standardisation of terminology in lower urinary tract function: report from the standardisation sub-committee of the International Continence Society. *Urology* 2003;61:37–49. doi: 10.1016/s0090-4295(02)02243-4
- Marinkovic SP, Stanton SL. Incontinence and voiding difficulties associated with prolapse. *J Urol* 2004;171:1021–8. doi: 10.1097/01.ju.0000111782.37383.e2
- Mouritsen L, Larsen JP. Symptoms, bother and POPQ in women referred with pelvic organ prolapse. *Int Urogynecol J Pelvic Floor Dysfunct* 2003;14:122–7. doi: 10.1007/s00192-002-1024-1
- Ellerkmann RM, Cundiff GW, Melick CF, Nihira MA, Leffler K, Bent AE. Correlation of symptoms with location and severity of pelvic organ prolapse. *Am J Obstet Gynecol* 2001;185:1332–8. doi: 10.1067/mob.2001.119078
- Chung CP, Miskimins R, Kuehl TJ, Yandell PM, Shull BL. Permanent suture used in uterosacral ligament suspension offers better anatomical support than delayed absorbable suture. *Int Urogynecol J* 2012;23:223–7. doi: 10.1007/s00192-011-1556-3
- Bradley MS, Bickhaus JA, Amundsen CL, Newcomb LK, Truong T, Weidner AC, et al. Vaginal uterosacral ligament suspension: a retrospective cohort of absorbable and permanent suture groups. *Female Pelvic Med Reconstr Surg* 2018;24:207–12. doi: 10.1097/SPV.0000000000000451
- Tagliaferri V, Ruggieri S, Tacaliti C, Gentile C, Didonna T, D'asta M, et al. Comparison of absorbable and permanent sutures for laparoscopic sacrocervicopexy: a randomized controlled trial. *Acta Obstet Gynecol Scand* 2021;100:347–52. doi: 10.1111/aogs.13997
- Powell CR, Tachibana I, Eckrich B, Rothenberg J, Hathaway J. Securing mesh with delayed absorbable suture does not increase risk of prolapse recurrence after robotic sacral colpopexy. *J Endourology* 2021;35:944–9. doi: 10.1089/end.2018.0029
- Linder BJ, Anand M, Klingele CJ, Trabuco EC, Gebhart JB, Occhino JA. Outcomes of robotic sacrocolpopexy using only absorbable suture for mesh fixation. *Female Pelvic Med Reconstr Surg* 2017;23:13–6. doi: 10.1097/SPV.0000000000000326
- Rubino SR, Popiel P, Gorgy M, Drugge E, Grimes CL. Pattern of suture use for pelvic organ prolapse surgery. *Am J Obstet Gynecol* 2022;226:1331–2. doi: 10.1016/j.ajog.2021.12.106
- Lee J, Oh S, Jeon MJ. Suture complication rates and surgical outcomes according to the nonabsorbable suture materials used in vaginal uterosacral ligament suspension: polyester versus polypropylene. *J Minimally Invasive Gynecol* 2021;28:1503–7. doi: 10.1016/j.jmig.2020.12.008
- Wallace BC, Trikalinos TA, Lau J, Brodley C, Schmid CH. Semi-automated screening of biomedical citations for systematic reviews. *BMC Bioinformatics* 2010;11:55. doi: 10.1186/1471-2105-11-55
- Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, et al. Grading quality of evidence and strength of recommendations. *BMJ* 2004;328:1490. doi: 10.1136/bmj.328.7454.1490
- Higgins JP, Altman DG. Assessing risk of bias in included studies. In: Higgins JP, Green S, editors. *Cochrane handbook for systematic reviews of interventions*. John Wiley & Sons, Ltd; 2008.
- Wells G, Wells G, Shea B, Shea B, O'Connell D, Peterson J, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. *Ottawa Hospital*; 2014.
- Nyaga VN, Arbyn M, Aerts M. Metaprop: a Stata command to perform meta-analysis of binomial data. *Arch Public Health* 2014;72:39. doi: 10.1186/2049-3258-72-39
- Kowalski JT, Genadry R, Ten Eyck P, Bradley CS. A randomized controlled trial of permanent vs absorbable suture for uterosacral ligament suspension. *Int Urogynecol J* 2021;32:785–90. doi: 10.1007/s00192-020-04244-1
- Silveira SdRB, Auge AP, Jarmy-Dibella ZI, Margarido PF, Carrao S, Alves Rodrigues C, et al. A multicenter, randomized trial comparing pelvic organ prolapse surgical treatment with native tissue and synthetic mesh: a 5-year follow-up study. *Neurourol Urodyn* 2020;39:1002–11. doi: 10.1002/nau.24323
- Goldberg RP, Tomezsko JE, Winkler HA, Koduri S, Culligan PJ, Sand PK. Anterior or posterior sacrospinous vaginal vault suspension: long-term anatomic and functional evaluation. *Obstet Gynecol* 2001;98:199–204. doi: 10.1016/s0029-7844(01)01446-6
- Mothes AR, Wanzke L, Radosa MP, Runnebaum IB. Bilateral minimal tension sacrospinous fixation in pelvic organ prolapse: an observational study. *Eur J Obstet Gynecol Reprod Biol* 2015;188:1–5. doi: 10.1016/j.ejogrb.2015.02.022
- Leone Roberti Maggiore U, Alessandri F, Remorgida V, Venturini PL, Ferrero S. Vaginal sacrospinous colpopexy using the Capiro suture-capturing device versus traditional technique: feasibility and outcome. *Arch Gynecol Obstet* 2013;287:267–74. doi: 10.1007/s00404-012-2540-y

23. Peng P, Zhu L, Lang J-H, Wang W-Y, Shi H-H. Unilateral sacrospinous ligament fixation for treatment of genital prolapse. *Chin Med J* 2010;123:1995–8. doi: 10.3760/cma.j.issn.0366-6999.2010.15.008
24. Astepe BS, Karsli A, Köleli I, Aksakal OS, Terzi H, Kale A. Intermediate-term outcomes of laparoscopic pectopexy and vaginal sacrospinous fixation: a comparative study. *Int Braz J Urol* 2019;45:999–1007. doi: 10.1590/S1677-5538.IBJU.2019.0103
25. Salvat J, Slamani L, Vincent-Genod A, Walker D. Sacrospinous ligament fixation by palpation: variation of the Richter procedure. *Eur J Obstet Gynecol Reprod Biol* 1996;68:199–203. doi: 10.1016/0301-2115(96)02470-0
26. Shastri SS, Darawade SP, Yedage NR. Transvaginal sacrospinous ligament fixation: revisited in modern gynecology. *J South Asian Fed Obstet Gynaecol* 2020;12:155–8. doi:10.5005/jp-journals-10006-1782
27. Wheeler TL, Gerten KA, Richter HE, Duke AG, Varner RE. Outcomes of vaginal vault prolapse repair with a high uterosacral suspension procedure utilizing bilateral single sutures. *Int Urogynecol J* 2007;18:1207–13. doi: 10.1007/s00192-007-0305-0
28. Nager CW, Visco AG, Richter HE, Rardin CR, Komesu Y, Harvie HS, et al. Effect of sacrospinous hysteropexy with graft vs vaginal hysterectomy with uterosacral ligament suspension on treatment failure in women with uterovaginal prolapse: 5-year results of a randomized clinical trial. *Am J Obstet Gynecol* 2021;225:153.e1–31. doi: 10.1016/j.ajog.2021.03.012
29. Barber MD, Visco AG, Weidner AC, Amundsen CL, Bump RC. Bilateral uterosacral ligament vaginal vault suspension with site-specific endopelvic fascia defect repair for treatment of pelvic organ prolapse. *Am J Obstet Gynecol* 2000;183:1402–11. doi: 10.1067/mob.2000.111298
30. Mowat A, Wong V, Goh J, Krause H, Pelecanos A, Higgs P. A descriptive study on the efficacy and complications of the Capio (Boston Scientific) suturing device for sacrospinous ligament fixation. *Aust N Z J Obstet Gynaecol* 2018;58:119–24. doi: 10.1111/ajo.12720
31. Dungal G, Poudel R, Shrestha R, Karki A, Pradhan HK, Bhattachan K, et al. Outcome of sacrospinous ligament fixation of the vault during repair of pelvic organ prolapse. *J Nepal Health Res Counc* 2018;16:321–4. doi:10.3126/JNHRC.V16I3.21431
32. Greisen S, Axelsen SM, Bek KM, Guldberg R, Glavind-Kristensen M. Fast track sacrospinous ligament fixation: subjective and objective outcomes at 6 months. *BMC Womens Health* 2021;21:154. doi: 10.1186/s12905-021-01309-1
33. Jeffery ST, Doumouchtsis SK, Franco AVM, Fynes MM. High uterosacral ligament vault suspension at vaginal hysterectomy: objective and subjective outcomes of a modified technique. *J Obstet Gynaecol Res* 2009;35:539–44. doi: 10.1111/j.1447-0756.2008.00984.x
34. Wong MJ, Rezvan A, Bhatia NN, Yazdany T. Uterosacral ligament vaginal vault suspension using delayed absorbable monofilament suture. *Int Urogynecol J* 2011;22:1389–94. doi: 10.1007/s00192-011-1470-8
35. Spelzini F, Frigerio M, Manodoro S, Interdonato ML, Cesana MC, Verri D, et al. Modified McCall culdoplasty versus Shull suspension in pelvic prolapse primary repair: a retrospective study. *Int Urogynecol J* 2017;28:65–71. doi: 10.1007/s00192-016-3016-6
36. Schiavi MC, Perniola G, Di Donato V, Visentin VS, Vena F, Di Pinto A, et al. Severe pelvic organ prolapse treated by vaginal native tissue repair: long-term analysis of outcomes in 146 patients. *Arch Gynecol Obstet* 2017;295:917–22. doi: 10.1007/s00404-017-4307-y
37. Rappa C, Saccone G. Recurrence of vaginal prolapse after total vaginal hysterectomy with concurrent vaginal uterosacral ligament suspension: comparison between normal-weight and overweight women. *Am J Obstet Gynecol* 2016;215:601.e1–4. doi: 10.1016/j.ajog.2016.06.022
38. Chill HH, Navon I, Reuveni-Salzman A, Cohen A, Dick A, Shveiky D. Vaginal colposuspension using the uphold lite mesh system versus transvaginal hysterectomy with uterosacral ligament suspension for treatment of apical prolapse: a comparative study. *J Minimally Invasive Gynecol* 2021;28:1759–64. doi: 10.1016/j.jmig.2021.03.002
39. Matthews CA, Geller EJ, Henley BR, Kenton K, Myers EM, Dieter AA, et al. Permanent compared with absorbable suture for vaginal mesh fixation during total hysterectomy and sacrocolpopexy: a randomized controlled trial. *Obstet Gynecol* 2020;136:355–64. doi: 10.1097/AOG.0000000000003884
40. Tan-Kim J, Menefee SA, Lippmann Q, Lukacz ES, Luber KM, Nager CW. A pilot study comparing anatomic failure after sacrocolpopexy with absorbable or permanent sutures for vaginal mesh attachment. *Permanente J* 2014;18:40–4. doi: 10.7812/TPP/14-022
41. Culligan PJ, Salamon C, Priestley JL, Shariati A. Porcine dermis compared with polypropylene mesh for laparoscopic sacrocolpopexy: a randomized controlled trial. *Obstet Gynecol* 2013;121:143–51. doi: 10.1097/aog.0b013e31827558dc
42. Elliott DS, Krambeck AE, Chow GK. Long-term results of robotic assisted laparoscopic sacrocolpopexy for the treatment of high grade vaginal vault prolapse. *J Urol* 2006;176:655–9. doi: 10.1016/j.juro.2006.03.040
43. Ross JW, Preston M. Laparoscopic sacrocolpopexy for severe vaginal vault prolapse: five-year outcome. *J Minimally Invasive Gynecol* 2005;12:221–6. doi: 10.1016/j.jmig.2005.03.017
44. Davila HH, Abdelhameed S, Malave-Huertas D, Bigay FF, Crawford K, Friedenstab A, et al. Ultrasonography and robotic-assisted laparoscopic sacrocervicopexy with pubocervical fascia reconstruction: comparison with standard technique. *J Robotic Surg* 2020;14:759–66. doi: 10.1007/s11701-020-01051-0
45. Culligan PJ, Blackwell L, Goldsmith LJ, Graham CA, Rogers A, Heit MH. A randomized controlled trial comparing fascia lata and synthetic mesh for sacral colpopexy. *Obstet Gynecol* 2005;106:29–37. doi: 10.1097/01.AOG.0000165824.62167.c1
46. Gracia M, Perelló M, Bataller E, Espuña M, Parellada M, Genís D, et al. Comparison between laparoscopic sacral hysteropexy and subtotal hysterectomy plus cervicopexy in pelvic organ prolapse: a pilot study. *Neurourol Urodyn* 2015;34:654–8. doi: 10.1002/nau.22641
47. Nosti PA, Carter CM, Sokol AI, Tefera E, Iglesia CB, Park AJ, et al. Transvaginal versus transabdominal placement of synthetic mesh at time of sacrocolpopexy. *Female Pelvic Med Reconstr Surg* 2016;22:151–5. doi: 10.1097/SPV.0000000000000222
48. Benson AD, Kramer BA, Wayment RO, Schwartz BF. Supracervical robotic-assisted laparoscopic sacrocolpopexy for pelvic organ prolapse. *JLS* 2010;14:525–30. doi: 10.4293/108680810X12924466008006
49. Culligan PJ, Lewis C, Priestley J, Mushonga N. Long-term outcomes of robotic-assisted laparoscopic sacrocolpopexy using lightweight Y-mesh. *Female Pelvic Med Reconstr Surg* 2020;26:202–6. doi: 10.1097/SPV.0000000000000788
50. Salamon CG, Culligan PJ. Subjective and objective outcomes 1 year after robotic-assisted laparoscopic sacrocolpopexy. *J Robotic Surg* 2013;7:35–8. doi: 10.1007/s11701-012-0337-4
51. Salamon CG, Lewis C, Priestley J, Gurshumov E, Culligan PJ. Prospective study of an ultra-lightweight polypropylene Y mesh

for robotic sacrocolpopexy. *Int Urogynecol J* 2013;24:1371–5. doi: 10.1007/s00192-012-2021-7

52. Tate SB, Blackwell L, Lorenz DJ, Steptoe MM, Culligan PJ. Randomized trial of fascia lata and polypropylene mesh for abdominal sacrocolpopexy: 5-year follow-up. *Int Urogynecol J* 2011;22:137–43. doi: 10.1007/s00192-010-1249-3
53. Martin LA, Calixte R, Finamore PS. Reoperation after robotic and vaginal mesh reconstructive surgery: a retrospective cohort study. *Female Pelvic Med Reconstr Surg* 2015;21:315–8. doi: 10.1097/SPV.0000000000000194
54. Sato H, Abe H, Ikeda A, Miyagawa T, Sato K. Complications and clinical outcomes of laparoscopic sacrocolpopexy for pelvic organ prolapse. *J Obstet Gynaecol* 2021;41:128–32. doi: 10.1080/01443615.2020.1724914
55. Kenton K, Mueller ER, Tarney C, Bresee C, Anger JT. One-year outcomes after minimally invasive sacrocolpopexy. *Female Pelvic Med Reconstr Surg* 2016;22:382–4. doi: 10.1097/SPV.0000000000000300
56. Belsante M, Murray S, Dillon B, Zimmern P. Mid term outcome of robotic mesh sacrocolpopexy. *Can J Urol* 2013;20:6656–61.
57. Elliott DS, Siddiqui SA, Chow GK. Assessment of the durability of robot-assisted laparoscopic sacrocolpopexy for treatment of vaginal vault prolapse. *J Robotic Surg* 2007;1:163–8. doi: 10.1007/s11701-007-0028-8
58. Mueller MG, Jacobs KM, Mueller ER, Abernethy MG, Kenton KS. Outcomes in 450 women after minimally invasive abdominal sacrocolpopexy for pelvic organ prolapse. *Female Pelvic Med Reconstr Surg* 2016;22:267–71. doi: 10.1097/SPV.0000000000000269
59. Van den Akker CM, Klerkx WM, Kluivers KB, van Eijndhoven HWF, Withagen MJJ, Scholten PC. Long-term safety, objective and subjective outcomes of laparoscopic sacrocolpopexy without peritoneal closure. *Int Urogynecol J* 2020;31:1593–600. doi: 10.1007/s00192-019-04020-w
60. Giannini A, Russo E, Misasi G, Falcone M, Caretto M, Morganti R, et al. Technical features, perioperative and anatomical outcomes of a standardized suturing pattern for robotic sacrocolpopexy. *Int Urogynecol J* 2022;33:3085–92. doi: 10.1007/s00192-021-05049-6
61. Bazzi AA, Osmundsen BC, Hagglund KH, Aslam MF. Anatomical outcomes based on suturing technique during vaginal mesh attachment in robotic sacrocolpopexy. *Female Pelvic Med Reconstr Surg* 2019;25:105–8. doi: 10.1097/SPV.0000000000000654
62. Morciano A, Marzo G, Caliendo D, Campagna G, Panico G, Alcaino S, et al. Laparoscopic sacral colpopexy and a new approach to mesh fixation: a randomized clinical trial. *Arch Gynecol Obstet* 2018;298:939–44. doi: 10.1007/s00404-018-4916-0
63. Berger AA, Tan-Kim J, Menefee SA. Anchor vs suture for the attachment of vaginal mesh in a robotic-assisted sacrocolpopexy: a randomized clinical trial. *Am J Obstet Gynecol* 2020;223:258.e1–8. doi: 10.1016/j.ajog.2020.05.018
64. Cvach K, Geoffrion R, Cundiff GW. Abdominal sacral hysterectomy: a pilot study comparing sacral hysterectomy to sacral colpopexy with hysterectomy. *Female Pelvic Med Reconstr Surg* 2012;18:286–90. doi: 10.1097/SPV.0b013e3182673772

65. Tan-Kim J, Nager CW, Grimes CL, Lubner KM, Lukacz ES, Brown HW, et al. A randomized trial of vaginal mesh attachment techniques for minimally invasive sacrocolpopexy. *Int Urogynecol J* 2015;26:649–56. doi: 10.1007/s00192-014-2566-8
66. Gilleran JP, Zimmern P. Abdominal mesh sacrocolpopexy for recurrent triple-compartment pelvic organ prolapse. *BJU Int* 2009;103:1090–4. doi: 10.1111/j.1464-410X.2008.08296.x
67. Liu J, Kohn J, Wu C, Guan Z, Guan X. Short-term outcomes of non-robotic single-incision laparoscopic sacrocolpopexy: a surgical technique. *J Minimally Invasive Gynecol* 2020;27:721–7. doi: 10.1016/j.jmig.2019.05.015
68. Borahay MA, Oge T, Walsh TM, Patel PR, Rodriguez AM, Kilic GS. Outcomes of robotic sacrocolpopexy using barbed delayed absorbable sutures. *J Minimally Invasive Gynecol* 2014;21:412–6. doi: 10.1016/j.jmig.2013.11.002
69. Shepherd JP, Higdon HL, Stanford EJ, Mattox TF. Effect of suture selection on the rate of suture or mesh erosion and surgery failure in abdominal sacrocolpopexy. *Female Pelvic Med Reconstr Surg* 2010;16:229–33. doi: 10.1097/SPV.0b013e3181d683a3
70. Stubbs JT. Short-term results of robotic sacrocolpopexy using the Quill SRS bi-directional polydioxanone (PDO) suture. *J Robotic Surg* 2011;5:259–65. doi: 10.1007/s11701-011-0267-6
71. Shekhar S, Goyal S, Goel M, Sharma C, Jhirwal M. Medium-term anatomical and functional results of isolated laparoscopic sacrocolpopexy for female pelvic organ prolapse during the early learning curve. *J Obstet Gynecol India* 2021;71:91–4. doi: 10.1007/s13224-020-01373-x
72. Balsamo R, Illiano E, Zucchi A, Natale F, Carbone A, Sio MD, et al. Sacrocolpopexy with polyvinylidene fluoride mesh for pelvic organ prolapse: mid term comparative outcomes with polypropylene mesh. *Eur J Obstet Gynecol Reprod Biol* 2018;220:74–8. doi: 10.1016/j.ejogrb.2017.11.018
73. Kallidonis P, Al-Aown A, Vasilas M, Kyriazis I, Panagopoulos V, Fligou F, et al. Laparoscopic sacrocolpopexy using barbed sutures for mesh fixation and peritoneal closure: a safe option to reduce operational times. *Urol Ann* 2017;9:159–65. doi: 10.4103/UA.UA_161_16
74. Liu J, Bardawil E, Zurawin RK, Wu J, Fu H, Orejuela F, Guan X. Robotic single-site sacrocolpopexy with retroperitoneal tunneling. *JSLs* 2018;22:e2018.00009.00009. doi: 10.4293/JSLs.2018.00009
75. Reisenauer C, Andress J, Schoenfish B, Huebner M, Brucker SY, Lippkowski A, et al. Absorbable versus non-absorbable sutures for vaginal mesh attachment during sacrocolpopexy: a randomized controlled trial. *Int Urogynecol J* 2022;33:411–9. doi: 10.1007/s00192-021-04853-4
76. Peng L, Liu Y-H, He S-X, Di X-P, Shen H, Luo D-Y. Is absorbable suture superior to permanent suture for uterosacral ligament suspension? *Neurourol Urodyn* 2020;39:1958–65. doi: 10.1002/nau.24434

PEER REVIEW HISTORY

Received July 11, 2022. Received in revised form September 2, 2022. Accepted September 8, 2022. Peer reviews and author correspondence are available at <http://links.lww.com/AOG/C975>.